

A generalization of the Tutte polynomials ^{*}

Tsuyoshi Miezuki [†], Manabu Oura [‡],
Tadashi Sakuma [§] and Hidehiro Shinohara [¶]

Abstract

In this paper, we introduce the concept of the Tutte polynomials of genus g and discuss some of its properties. We note that the Tutte polynomials of genus one are well-known Tutte polynomials. The Tutte polynomials are matroid invariants, and we claim that the Tutte polynomials of genus g are also matroid invariants. The main result of this paper and the forthcoming paper declares that the Tutte polynomials of genus g are complete matroid invariants.

Key Words and Phrases. matroid, Tutte polynomial.

2010 *Mathematics Subject Classification.* Primary 05B35.

1 Introduction

This is an announcement paper.

Let E be a set. A *matroid* M on $E = E(M)$ is a pair (E, \mathcal{I}) , where \mathcal{I} is a non-empty family of subsets of E with the properties

$$\left\{ \begin{array}{ll} \text{(i)} & \text{if } I \in \mathcal{I} \text{ and } J \subset I, \text{ then } J \in \mathcal{I}; \\ \text{(ii)} & \text{if } I_1, I_2 \in \mathcal{I} \text{ and } |I_1| < |I_2|, \\ & \text{then there exists } e \in I_2 \setminus I_1 \\ & \text{such that } I_1 \cup \{e\} \in \mathcal{I}. \end{array} \right.$$

^{*}This work was supported by JSPS KAKENHI(18K03217, 17K05164, 18K03388).

[†]Faculty of Education, Ryukyu University, Okinawa, 903–0213, Japan

[‡]Graduate School of Natural Science and Technology, Kanazawa University, Ishikawa 920–1192, Japan

[§]Faculty of Science, Yamagata University, Yamagata 990–8560, Japan

[¶]Institute for Excellence in Higher Education, Tohoku University, Miyagi 980–8576, Japan

Each element of the set \mathcal{I} is called an *independent set*. A matroid (E, \mathcal{I}) is *isomorphic* to another matroid (E', \mathcal{I}') if there is a bijection φ from E to E' such that $\varphi(I) \in \mathcal{I}'$ holds for each member $I \in \mathcal{I}$, and $\varphi^{-1}(I') \in \mathcal{I}$ holds for each member $I' \in \mathcal{I}'$.

It follows from the second axiom that all maximal independent sets in a matroid M take the same cardinality, called the *rank* of M . These maximal independent sets $\mathcal{B}(M)$ are called the *bases* of M . The *rank* $\rho(A)$ of an arbitrary subset A of E is the cardinality of the largest independent set contained in A .

We provide examples below.

Example 1.1. Let A be a $k \times n$ matrix over a finite field \mathbb{F}_q . This offers a matroid M on the set

$$E = \{z \in \mathbb{Z} \mid 1 \leq z \leq n\}$$

in which a set I is independent if and only if the family of columns of A whose indices belong to I is linearly independent. Such a matroid is called a vector matroid.

Example 1.2. Let

$$E = \{z \in \mathbb{Z} \mid 1 \leq z \leq n\}$$

and r a natural number. We define a matroid on E by taking every r -element subset of E to be a basis. This is known as the uniform matroid $U_{r,n}$.

The classification of the matroids is one of the most important problems in the theory of matroids. One tool to classify the matroids is the Tutte polynomials. Let M be a matroid on the set E having a rank function ρ . The Tutte polynomial of M is defined as follows [2, 3, 4]:

$$\begin{aligned} T(M) &:= T(M; x, y) \\ &:= \sum_{A \subseteq E} (x-1)^{\rho(E)-\rho(A)} (y-1)^{|A|-\rho(A)}. \end{aligned}$$

For example, the Tutte polynomial of the uniform matroid $U_{r,n}$ is

$$\begin{aligned} T(U_{r,n}; x, y) &= \sum_{i=0}^r \binom{n}{i} (x-1)^{r-i} + \sum_{i=r+1}^n \binom{n}{i} (y-1)^{i-r}. \end{aligned}$$

It is easy to demonstrate that $T(M; x, y)$ is a matroid invariant. Two matroids are called *T-equivalent* if their Tutte polynomials are equivalent.

It is well known that there exist two inequivalent matroids, which are *T-equivalent* [5, p. 269]. We provide more examples below. Let

$$E := \{z \in \mathbb{Z} \mid 1 \leq z \leq n\}.$$

Let us define the subsets X_1, X_2, X_3 of E as follows:

$$\left\{ \begin{array}{l} X_1 := \{z \in \mathbb{Z} \mid 1 \leq z \leq r\}; \\ X_2 := \{z \in \mathbb{Z} \mid r + 1 \leq z \leq 2r\}; \\ X_3 := \{z \in \mathbb{Z} \mid r \leq z \leq 2r - 1\}. \end{array} \right.$$

Let $R_{r,n}$ denote the matroid on E such that

$$\mathcal{B}(R_{r,n}) = \mathcal{B}(U_{r,n}) \setminus \{X_1, X_2\},$$

$Q_{r,2n}$ denote the matroid on E such that

$$\mathcal{B}(Q_{r,n}) = \mathcal{B}(U_{r,n}) \setminus \{X_1, X_3\}.$$

Then, for $2r \leq n$, $r \geq 3$, $R_{r,n}$ and $Q_{r,n}$ are matroids. Both matroids have exactly two dependent sets of size r , namely $\{X_1, X_2\}$ of $R_{r,n}$ and $\{X_1, X_3\}$ of $Q_{r,n}$. Therefore, if $R_{r,n}$ and $Q_{r,n}$ are isomorphic, there exists φ such that

$$\varphi(\{X_1, X_2\}) = \{X_1, X_3\}.$$

This is a contradiction since φ is bijective, and we know that $R_{r,n}$ and $Q_{r,n}$ are non-isomorphic matroids.

On the other hand,

$$T(R_{r,n}) = T(Q_{r,n}).$$

In fact, the difference between

$$T(U_{r,n}) - T(R_{r,n})$$

and

$$T(U_{r,n}) - T(Q_{r,n})$$

are zero since $R_{r,n}$ and $Q_{r,n}$ are obtained from $U_{r,n}$ after deleting the two maximal independent sets.

This gives rise to a natural question: is there a generalization of the Tutte polynomial which identifies such T -equivalent but inequivalent matroids? This paper aims to provide a candidate generalization that answers this. In Section 2, we provide the concept of the Tutte polynomial of genus g . In Section 3, we provide the main results. The details of the proofs will be presented in our forthcoming paper [1].

2 Tutte polynomials of genus g

We now present the concept of the Tutte polynomial of genus g .

Definition 2.1. Let $M := (E, \mathcal{I})$ be a matroid. Let

$$\Lambda_1 := \{z \in \mathbb{Z} \mid 1 \leq z \leq g\}$$

and let

$$\Lambda_2 := \binom{\Lambda_1}{2}.$$

For every element $\lambda \in \Lambda_2$, let us denote

$$A_{\cap(\lambda)} := \cap_{i \in \lambda} A_i \text{ and } A_{\cup(\lambda)} := \cup_{i \in \lambda} A_i.$$

Then, the genus g of the Tutte polynomial $T^{(g)}(M)$ of the matroid M will be defined as follows:

$$\begin{aligned} T^{(g)}(M) &:= T^{(g)}(M; x_{\lambda_1}, y_{\lambda_1}, x_{\cap \lambda_2}, y_{\cap \lambda_2}, \\ &\quad x_{\cup \lambda_2}, y_{\cup \lambda_2} : \lambda_1 \in \Lambda_1, \lambda_2 \in \Lambda_2) \\ &:= \sum_{A_1, \dots, A_n \subseteq E} \prod_{\lambda \in \Lambda_1} (x_\lambda - 1)^{\rho(M) - \rho(A_\lambda)} \\ &\quad (y_\lambda - 1)^{|A_\lambda| - \rho(A_\lambda)} \\ &\quad \prod_{\lambda \in \Lambda_2} (x_{\cap(\lambda)} - 1)^{\rho(M) - \rho(A_{\cap(\lambda)})} \\ &\quad (y_{\cap(\lambda)} - 1)^{|A_{\cap(\lambda)}| - \rho(A_{\cap(\lambda)})} \\ &\quad \prod_{\lambda \in \Lambda_2} (x_{\cup(\lambda)} - 1)^{\rho(M) - \rho(A_{\cup(\lambda)})} \\ &\quad (y_{\cup(\lambda)} - 1)^{|A_{\cup(\lambda)}| - \rho(A_{\cup(\lambda)})}. \end{aligned}$$

It is easy to demonstrate that $T^{(g)}(M)$ is matroid invariant and if two matroids have the same Tutte polynomial for genus g , we call them $T^{(g)}$ -equivalent. For example, the genus 2 for the Tutte polynomial $T^{(2)}(M)$ of the matroid M is as follows:

$$\begin{aligned}
& T^{(2)}(M; x_1, x_2, y_1, y_2, x_{\{1,2\}}, y_{\{1,2\}}, x_{\cup\{1,2\}}, y_{\cup\{1,2\}}) \\
&= \sum_{A_1, A_2 \subset E} (x_1 - 1)^{\rho(E) - \rho(A_1)} \\
&\quad (x_2 - 1)^{\rho(E) - \rho(A_2)} \\
&\quad (y_1 - 1)^{|A_1| - \rho(A_1)} \\
&\quad (y_2 - 1)^{|A_2| - \rho(A_2)} \\
&\quad (x_{\{1,2\}} - 1)^{\rho(E) - \rho(A_1 \cap A_2)} \\
&\quad (y_{\{1,2\}} - 1)^{|A_1 \cap A_2| - \rho(A_1 \cap A_2)} \\
&\quad (x_{\cup\{1,2\}} - 1)^{\rho(E) - \rho(A_1 \cup A_2)} \\
&\quad (y_{\cup\{1,2\}} - 1)^{|A_1 \cup A_2| - \rho(A_1 \cup A_2)}.
\end{aligned}$$

We remark that for $g \in \mathbb{N}_{\geq 2}$, the specialization of $T^{(g)}(M)$ is $T^{(g-1)}(M)$. For example,

$$T^{(2)}(M; x_1, 2, y_1, 2, 2, 2, 2) = 2^{|E|} T^{(1)}(M; x_1, y_1).$$

Therefore, if

$$T^{(g)}(M) = T^{(g)}(M')$$

then

$$T^{(i)}(M) = T^{(i)}(M'),$$

for all $1 \leq i \leq g$.

3 Main results

The main result of the present paper is as follows:

Theorem 3.1. *1. The Tutte polynomial of genus g $\{T^{(g)}\}_{g=1}^{\infty}$ is a complete invariant for matroids.*

2. For $2r \leq n$, $r \geq 3$,

$$T^{(2)}(R_{r,n}) \neq T^{(2)}(Q_{r,n}).$$

3. Let

$$E := \{z \in \mathbb{Z} \mid 0 \leq z \leq 4n - 1\}.$$

with $n \geq 3$. Let us define the subsets Y_1, Y_2 of 2^E as follows:

$$\begin{aligned} Y_1 := & \{\{0, 1, 2\}, \{2, 3, 4\}, \dots, \\ & \{2n - 4, 2n - 3, 2n - 2\}, \\ & \{2n - 2, 2n - 1, 0\}, \\ & \{2n, 2n + 1, 2n + 2\}, \\ & \{2n + 2, 2n + 3, 2n + 4\}, \dots, \\ & \{4n - 4, 4n - 3, 4n - 2\}, \\ & \{4n - 2, 4n - 1, 2n\}\}, \\ Y_2 := & \{\{0, 1, 2\}, \{2, 3, 4\}, \dots, \\ & \{2n - 4, 2n - 3, 2n - 2\}, \\ & \{2n - 2, 2n - 1, 2n\}, \\ & \{2n, 2n + 1, 2n + 2\}, \\ & \{2n + 2, 2n + 3, 2n + 4\}, \dots, \\ & \{4n - 4, 4n - 3, 4n - 2\}, \\ & \{4n - 2, 4n - 1, 0\}\}. \end{aligned}$$

Let S_{4n} denote the independence system on E such that

$$\mathcal{B}(S_{4n}) = \mathcal{B}(U_{3,4n}) \setminus Y_1,$$

S'_{4n} denote the independence system on E such that

$$\mathcal{B}(S'_{4n}) = \mathcal{B}(U_{3,4n}) \setminus Y_2.$$

Then, S_{4n} and S'_{4n} are matroids. Let

$$m_1 = \left\lfloor \frac{-1 + \sqrt{1 + 8n}}{2} \right\rfloor \quad \text{and} \quad m_2 = 2\lceil \sqrt{n} \rceil.$$

We have

$$\begin{cases} T^{(m_1)}(S_{4n}) = T^{(m_1)}(S'_{4n}); \\ T^{(m_2)}(S_{4n}) \neq T^{(m_2)}(S'_{4n}). \end{cases}$$

In particular, for a matroid M , $T^{(|\mathcal{B}(M)|)}(M)$ determines M . For detailed explanation, see [1].

Acknowledgments

This work was supported by JSPS KAKENHI (18K03217, 17K05164, 18K03388).

References

- [1] T. Miezeki, M. Oura, T. Sakuma, H. Shinohara, The Tutte polynomials of genus g , in preparation.
- [2] W. T. Tutte, A ring in graph theory, *Proc. Cambridge Philos. Soc.*, 43:26–40, 1947.
- [3] W. T. Tutte, A contribution to the theory of chromatic polynomials, *Canadian J. Math.*, 6:80–91, 1954.
- [4] W. T. Tutte, On dichromatic polynomials, *J. Combinatorial Theory*, 2:301–320, 1967.
- [5] D. J. A. Welsh, Matroid Theory, *Academic Press, London*, 1976.