

AN EXAMPLE OF AN INFINITELY GENERATED GRADED RING MOTIVATED BY CODING THEORY

By MANABU OURA¹

Abstract

Let $\tilde{\mathfrak{H}}^{(g)}$ be the graded ring generated by the r -th higher weight enumerators of all codes of any length, $1 \leq r \leq g$. In this note we will prove that $\tilde{\mathfrak{H}}^{(g)}$ is infinitely generated.

1. *Preliminaries.* Let \mathbf{F}_q be the field of q elements. A code C of length n means a subspace of \mathbf{F}_q^n . For the details of coding theory and the weight enumerators we shall next define, we refer to [4] and its references.

Let C be a code of length n and $W_C^{(g)}(x_a : a \in \mathbf{F}_q^g)$ the g -th complete weight enumerator of the code C , that is,

$$W_C^{(g)}(x_a : a \in \mathbf{F}_q^g) = \sum_{v_1, \dots, v_g \in C} \prod_{a \in \mathbf{F}_q^g} x_a^{n_a(v_1, \dots, v_g)},$$

where $n_a(v_1, \dots, v_g)$ denotes the number of i such that $a = (v_{1i}, \dots, v_{gi})$. This is a homogeneous² polynomial of degree n . Direct computation shows

$$W_{C_1 \oplus C_2}^{(g)}(x_a : a \in \mathbf{F}_q^g) = W_{C_1}^{(g)}(x_a : a \in \mathbf{F}_q^g) W_{C_2}^{(g)}(x_a : a \in \mathbf{F}_q^g) \quad (1)$$

for codes C_1 and C_2 , where \oplus denotes the direct sum of codes.

We write $W_C^{(g)}(x, y)$ after transforming the variables of $W_C^{(g)}(x_a : a \in \mathbf{F}_q^g)$ as

$$x_a \rightsquigarrow \begin{cases} x & \text{if } a = 0 \in \mathbf{F}_q^g, \\ y & \text{otherwise.} \end{cases}$$

We inductively define the r -th weight enumerator by the identity

$$W_C^{(g)}(x, y) = \sum_{0 \leq r \leq g} [g]_r H_C^{(r)}(x, y), \quad (2)$$

¹This work is supported in part by KAKENHI (No. 14740081).

²Throughout this note we assume that each degree of x and y is 1, thus the degree of $x^i y^j$ is $i + j$.

where $[g]_0 = 1$ and $[g]_r = (q^g - q^{r-1})(q^g - q^{r-2}) \cdots (q^g - q)(q^g - 1)$ for $1 \leq r \leq g$. Compare with [1], [2].

2. Result. Let $\tilde{\mathfrak{H}}^{(g)}$ be the ring generated by the $H_C^{(r)}(x, y)$'s, $1 \leq r \leq g$, of all codes of any length. Our result in this note is

Theorem. The ring $\tilde{\mathfrak{H}}^{(g)}$ is infinitely generated.

In order to prove Theorem we need the following

Lemma. Let C be a code of length n . Then $x^n H_C^{(r)}(x, y)$ is contained in $\tilde{\mathfrak{H}}^{(g)}$ for any r , $1 \leq r \leq g$.

Proof. We prove this by induction on r . Let C be a code of length n . If $r = 1$, we have

$$W_{C \oplus C}^{(1)}(x, y) = x^{2n} + H_{C \oplus C}^{(1)}(x, y).$$

Using the identity (1), we have

$$2x^n H_C^{(1)}(x, y) = H_{C \oplus C}^{(1)}(x, y) - \left(H_C^{(1)}(x, y) \right)^2.$$

The right hand side of this formula, thus $x^n H_C^{(1)}(x, y)$, lies in $\tilde{\mathfrak{H}}^{(g)}$.

Suppose that $r \geq 2$. Considering $C \oplus C$ instead of C in the identity (2), we have

$$\left(\sum_{i=0}^r [r]_i H_C^{(i)}(x, y) \right)^2 = \sum_{i=0}^r [r]_i H_{C \oplus C}^{(i)}(x, y),$$

or,

$$\begin{aligned} 2x^n H_C^{(r)}(x, y) &= \sum_{i=1}^r [r]_i H_{C \oplus C}^{(i)}(x, y) - \\ &\quad \left\{ \left(\sum_{i=1}^{r-1} [r]_i H_C^{(i)}(x, y) \right)^2 + \left([r]_r H_C^{(r)}(x, y) \right)^2 + \right. \\ &\quad \left. 2x^n \left(\sum_{i=1}^{r-1} [r]_i H_C^{(i)}(x, y) \right) + 2 \left(\sum_{i=1}^{r-1} [r]_i H_C^{(i)}(x, y) \right) [r]_r H_C^{(r)}(x, y) \right\}. \end{aligned}$$

Applying the induction hypothesis to the right hand side of this formula, we have that $x^n H_C^{(r)}(x, y)$ lies in $\tilde{\mathfrak{H}}^{(g)}$. This completes the proof of Lemma.

Proof of Theorem. If

$$\begin{aligned} f &= a_n x^n + a_{n-1} x^{n-1} y + \cdots + a_0 y^n, \\ a_n &= \cdots = a_{\ell-1} = 0, \quad a_\ell \neq 0, \end{aligned}$$

then we write $w(f) = \ell$. We put $w(0) = \infty$. For any code C of length n and any positive integer r , we have $w(H_C^{(r)}(x, y)) < n$. This fact will be used below.

Preparing this, we shall show that $\tilde{\mathfrak{H}}^{(g)}$ is infinitely generated. Assume that $\tilde{\mathfrak{H}}^{(g)}$ is finitely generated: $\tilde{\mathfrak{H}}^{(g)} = \mathbf{C}[H_{C_1}^{(i_1)}(x, y), \dots, H_{C_k}^{(i_k)}(x, y)]$. For a positive integer d , we shall denote by $\tilde{\mathfrak{H}}^{(g)}(d)$ the subring of $\tilde{\mathfrak{H}}^{(g)}$ generated by all elements of $\tilde{\mathfrak{H}}^{(g)}$ whose degrees are multiples of d . The degrees of the generators of $\tilde{\mathfrak{H}}^{(g)}$ may be different, however, certain subring of $\tilde{\mathfrak{H}}^{(g)}$ is able to be generated by the elements whose degrees are the same. More precisely there exists a positive integer r such that $\tilde{\mathfrak{H}}^{(g)}(r)$ can be generated by the F_1, \dots, F_m whose degrees (as homogeneous polynomials in $\mathbf{C}[x, y]$) are r (cf. [3], p.89 Lemma 3). Here we may take each F_i as a monomial of $H_{C_1}^{(i_1)}(x, y), \dots, H_{C_k}^{(i_k)}(x, y)$. Moreover we assume $w(F_1) \leq w(F_2) \leq \cdots \leq w(F_m)$. We remark that $w(F_m) < r$ because of the fact stated after the definition of $w(*)$. By Lemma, $x^r F_m$ belongs to $\tilde{\mathfrak{H}}^{(g)}(r)$ and can be written in the form

$$x^r F_m = \sum (\text{const.}) F_i F_j.$$

But this is impossible because of $w(x^r F_m) = r + w(F_m) > w(F_i F_j)$ for any i, j . Hence $\tilde{\mathfrak{H}}^{(g)}$ is infinitely generated.

DIVISION OF MATHEMATICS
SCHOOL OF MEDICINE
SAPPORO MEDICAL UNIVERSITY

REFERENCES

- [1] Dougherty, S. T., Gulliver, T. A., Oura, M., Higher weights and binary self-dual codes, to appear in *Discrete Applied Math*.
- [2] Dougherty, S. T., Gulliver, T. A., Oura, M., Higher Weights for Ternary and Quaternary Self-Dual Codes, preprint.
- [3] Igusa, J., Theta functions, *Grundlehren der Math. Wiss.* vol 194, Berlin Heidelberg New York: Springer (1972).
- [4] Nebe, G., Rains, E. M., Sloane, N. J. A., The invariants of the Clifford groups, *Designs, Codes and Cryptography*, **24** (2001), 99–122.